

1818 New York Avenue Suite 217 Washington, DC 20002

www.globalincusa.net

December 9, 2020

Prince Georges County Environmental Safety Office 13306 Old Marlboro Pike Upper Marlboro, MD 20772

Attention: Mr. Alex Baylor

RE: Indoor Air Quality Screening Report

Global Project Number: 20-064

School: Beacon Heights Elementary School

Dear Mr. Baylor,

On November 30, 2020, Global Inc.'s (GLOBAL) team of Industrial Hygienists under the supervision of Certified Industrial Hygienist, Dr. Channa Bambaradeniya, conducted an Indoor Air Quality Screening at Beacon Heights Elementary School located at 6929 Furman Parkway, Riverdale, MD.

Methodology

The IAQ evaluation included a visual assessment, sampling for non-viable mold spores in air, and measurement of comfort parameters (temperature, humidity, carbon dioxide, and carbon monoxide) in randomly selected representative locations within the building. GLOBAL's inspector conducted a walkthrough with Prince Georges County Public School (PGCPS) personnel present. Rooms were selected in a random manner throughout the building so as to prevent sampling bias.

During the visual assessment of representative locations, and when noted, GLOBAL documented those areas with suspected mold growth, water intrusions, and wet conditions that have the potential to lead to mold growth. GLOBAL also noted any unusual odors. At least one microbial air sample was collected for every 10,000 Square Feet (SF) of space in the building and the analytical results for the interior spaces were compared to an outdoor (ambient) sample collected on the same day.

Microbial samples (including a field blank for quality control) were delivered under strict chain-of-custody procedures were to Hayes Microbial Consulting - an AIHA EMPAT-certified laboratory in Midlothian, Virginia for analysis by microscopy. The sample chain-of-custody and laboratory report is attached.



1818 New York Avenue Suite 217 Washington, DC 20002 www.globalincusa.net

Observations

The general observations in the three indoor locations inspected are summarized in Table 1 below:

Table 1: Observations

Location	Observations					
Room 19	No issues					
Room 9	No issues					
Multi-purpose Room	No issues					

Comfort Parameter Measurements and Mold-in-Air Sample Results

The comfort parameter measurements and status of fungal ecology is summarized in Table 2.

Temperature

The American Society of Heating, Refrigerating and Air-Conditioning Engineers (ASHRAE) have published recommendations for year-round acceptable temperatures in Standard 55-2016 (*Thermal Environmental Conditions for Human Occupancy*). The winter comfort range is 68 to 75°F and the summer comfort range is 73 to 79°F. It is important to note that ASHRAE standards are intended as a suggested guideline as opposed to a regulation. All the indoor temperature readings were within the ASHRAE Standard.

Relative Humidity (RH)

Relative humidity is a key factor for mold growth. Mold has the potential of growing on suitable surfaces with humidity levels above 60%. ASHRAE standard 62.1-2013 (*Ventilation for Acceptable Indoor Air Quality*) recommends a maximum indoor relative humidity of 65% to preclude the likelihood of condensation on cool surfaces encouraging mold growth. All the indoor relative humidity readings were below the ASHRAE recommended level of 65%.

Carbon Monoxide

Carbon monoxide (CO) is a colorless and odorless gas that is produced by the incomplete combustion of carbon-containing fuels. Oil, gasoline, diesel fuels, wood, coke, and coal are the major sources of CO. All registered CO concentrations were below the EPA National Ambient Air Quality Standard (NAAQS) of 9 ppm.

Carbon Dioxide

Under conditions of maximum occupancy, ASHRAE Standard 62.1-2013, Appendix C, infers that the acceptable carbon dioxide upper limit is the prevailing outdoor carbon dioxide concentration



1818 New York Avenue Suite 217 Washington, DC 20002

www.globalincusa.net

plus 700 parts per million (ppm). On November 30, 2020, the outdoor (ambient) carbon dioxide concentration was approximately 420 ppm so indoor concentrations should not exceed approximately 1120 ppm (700 + 420). All indoor carbon dioxide measurements were within the ASHRAE standards.

Mold-in-Air Samples

There are no definitive regulations or standardized guidelines for addressing airborne mold in an indoor setting. If building systems (ventilation, envelope) are functioning properly, the indoor fungal ecology profile should be consistent with what is encountered outdoors and the spore concentrations should be below the ambient levels. Laboratory analytical results are attached at the end of this report.

Table 2: Air Quality Results

Sample Location Standards	Temp ⁰ F ASHRAE 68 to 75°F	RH% ASHRAE <65%	CO ppm NAAQS <9	CO2 ppm ASHRAE 1120	Normal Fungal Ecology?	
Ambient	66.5	78.1	0	420	-	
Room 19	68.4	55.5	0	511	Yes	
Room 9	68.6	53.7	0	411	Yes	
Multi-purpose	72.3	62.9	0	442	Yes	



1818 New York Avenue Suite 217 Washington, DC 20002

www.globalincusa.net

Conclusions and Recommendations

All comfort parameters measured were within the applicable Standards. No indoor air quality issues related to mold were found during the screening performed on November 30, 2020, and all mold samples were found to have a normal ecology for an indoor environment.

It has been our pleasure to conduct these IAQ Screening services for the Prince Georges County Public School system. If you have any questions, please feel free to contact us.

Regards,

Channa Bambaradeniya, Ph.D., CIH, CSP, CHMM

Certified Industrial Hygienist

Global, Inc.

Mobile: 443-691-0455



1818 New York Avenue Suite 217 Washington, DC 20002 www.globalincusa.net

ATTACHMENT I

Air Sample Analytical Results and Chain-Of-Custody Form





Analysis Report prepared for

Global, Inc.

1818 New York Ave. Suite 217 Washington, DC, 20002

Phone: (443) 691-0455

BB203
PGCPS Indoor Air Quality
Beacon Heights Elementary School

Collected: November 30, 2020 Received: December 1, 2020 Reported: December 1, 2020 We would like to thank you for trusting Hayes Microbial for your analytical needs! We received 4 samples by FedEx in good condition for this project on December 1st, 2020.

The results in this analysis pertain only to this job, collected on the stated date, and should not be used in the interpretation of any other job. This report may not be duplicated, except in full, without the written consent of Hayes Microbial Consulting, LLC..

This laboratory bears no responsibility for sample collection activities, analytical method limitations, or your use of the test results. Interpretation and use of test results are your responsibility. Any reference to health effects or interpretation of mold levels is strictly the opinion of Hayes Microbial. In no event, shall Hayes Microbial or any of its employees be liable for lost profits or any special, incidental or consequential damages arising out of the use of these test results.

Steve Hayes, BSMT(ASCP)
Laboratory Director

Hayes Microbial Consulting, LLC.



EPA Laboratory ID: VA01419



plan N. Hayes

Lab ID: #188863



DPH License: #PH-0198

1818 New York Ave. Suite 217 Washington, DC, 20002 (443) 691-0455

BB203

PGCPS Indoor Air Quality Beacon Heights Elementary School #20044834

Spore Trap SOP - HMC#101

Sample Number	e Number 1 BHES-1130-01		130-01	2 BHES-1130-02			3 BHES-1130-03			4 BHES-1130-04			
Sample Name	75.00 liter 13 spores/m ³ 2			75.00 liter 13 spores/m ³ 2 ND			Room 9 75.00 liter 13 spores/m ³ 2 ND			Multipurpose 75.00 liter			
Sample Volume													
Reporting Limit										13 spores/m³ 2 27/m³			
Background													
Fragments													
Organism	Raw Count	Count / m ³	% of Total	Raw Count	Count / m ³	% of Total	Raw Count	Count / m ³	% of Total	Raw Count	Count / m ³	% of Tota	
Alternaria							1	13	14.3%				
Ascospores	38	507	88.4%							25	333	78.1%	
Aspergillus Penicillium	3	40	7.0%										
Basidiospores	2	27	4.7%							1	13	3.1%	
Bipolaris Drechslera							1	13	14.3%	1	13	3.1%	
Chaetomium													
Cladosporium							4	53	57.1%	3	40	9.4%	
Curvularia													
Epicoccum										1	13	3.1%	
Fusarium													
Memnoniella													
Myxomycetes				2	27	100.0%	1	13	14.3%	1	13	3.1%	
Pithomyces													
Stachybotrys													
Stemphylium													
Torula													
Ulocladium													
Total	43	574	100%	2	27	100%	7	92	100%	32	425	100%	

Water Damage Indicator

Collected: Nov 30, 2020

Common Allergen

Slightly Higher than Baseline

Significantly Higher than Baseline

Ratio Abnormality

Received: Dec 1, 2020

Reported: Dec 1, 2020

Project Analyst:

Shareef Abdelgadir, MS <

12 - 01 - 2020

Date:

Reviewed By:

Steve Hayes, BSMT

Date:

12 - 01 - 2020

3005 East Boundary Terrace, Suite F. Midlothian, VA. 23112

(804) 562-3435

contact@hayesmicrobial.com

Page: 2 of 5

1818 New York Ave. Suite 217 Washington, DC, 20002 (443) 691-0455

BB203 PGCPS Indoor Air Quality Beacon Heights Elementary School

#20044834

Spore Trap Information

Reporting Limit	The Reporting Limit is the lowest number of spores that can be detected based on the total volume of the sample collected and the percentage of the slide that is counted. At Hayes Microbial, 100% of the slide is read so the LOD is based solely on the total volume. Raw spore counts that exceed 500 spores will be estimated.
Blanks	Results have not been corrected for field or laboratory blanks.
Background	The Background is the amount of debris that is present in the sample. This debris consists of skin cells, dirt, dust, pollen, drywall dust and other organic and non-organic matter. As the background density increases, the likelihood of spores, especially small spores such as those of Aspergillus and Penicillium may be obscured. The background is rated on a scale of 1 to 5 and each level is determined as follows:
	 NBD: No background detected due to possible pump or cassette malfunction. Recollect sample. (Field Blanks will display NBD) 1: <5% of field occluded. No spores will be uncountable. 2: 5-25% of field occluded. 3: 25-75% of field occluded. 4: 75-90% of field occluded. 5: >90% of field occluded. Suggested recollection of sample.
Fragments	Fragments are small pieces of fungal mycelium or spores. They are not identifiable as to type and when present in very large numbers, may indicate the presence of mold amplification.
Control Comparisons	There are no national standards for the numbers of fungal spores that may be present in the indoor environment. As a general rule and guideline that is widely accepted in the indoor air quality field, the numbers and types of spores that are present in the indoor environment should not exceed those that are present outdoors at any given time. There will always be some mold spores present in "normal" indoor environments. The purpose of sampling and counting spores is to help determine whether an abnormal condition exists within the indoor environment and if it does, to help pinpoint the area of contamination. Spore counts should not be used as the sole determining factor of mold contamination. There are many factors that can cause anomalies in the comparison of indoor and outdoor samples due to the dynamic nature of both of those environments.
Water Damage Indicator	Blue: These molds are commonly seen in conditions of prolonged water intrusion and usually indicate a problem.
Common Allergen	Green: Although all molds are potential allergens, these are the most common allergens that may be found indoors.
Slightly Higher than Baseline	Orange: The spore count is slightly higher than the outside count and may or may not indicate a source of contamination.
Significantly Higher than Baseline	Red: The spore count is significantly higher than the baseline count and probably indicates a source of contamination.
Ratio Abnormality	Violet: The types of spores found indoors should be similar to the ones that were identified in the baseline sample. Significant increases (more than 25%) in the ratio of a particular spore type may indicate the presence of abnormal levels of mold, even if the total number of spores of that type is lower in the indoor environment than it was outdoors.
Color Coding	Fungi that are present in indoor samples at levels lower than 200 per cubic meter are not color coded on the report, unless they are one of the water damage indicators.




1818 New York Ave. Suite 217 Washington, DC, 20002 (443) 691-0455

BB203 PGCPS Indoor Air Quality Beacon Heights Elementary School

#20044834

Organism Descriptions

Alternaria	Habitat: Effects:	Commonly found outdoors in soil and decaying plants. Indoors, it is commonly found on window sills and other horizontal surfaces. A common allergen and has been associated with hypersensitivity pneumonitis. Alternaria is capable of producing toxic metabolites which may be associated with disease in humans or animals. Occasionally an agent of onychomycosis, ulcerated cutaneous infection and chronic sinusitis, principally in the immunocompromised patient.
Ascospores	Habitat: Effects:	A large group consisting of more than 3000 species of fungi. Common plant pathogens and outdoor numbers become very high following rain. Most of the genera are indistinguishable by spore trap analysis and are combined on the report. Health affects are poorly studied, but many are likely to be allergenic.
Aspergillus Penicillium	Habitat:	The most common fungi isolated from the environment. Very common in soil and on decaying plant material. Are able to grow well indoors on a wide variety of substrates.
	Effects:	This group contains common allergens and many can cause hypersensitivity pneumonitis. They may cause extrinsic asthma, and many are opportunistic pathogens. Many species produce mycotoxins which may be associated with disease in humans and other animals. Toxin production is dependent on the species, the food source, competition with other organisms, and other environmental conditions.
Basidiospores	Habitat:	A common group of Fungi that includes the mushrooms and bracket fungi. They are saprophytes and plant pathogens. In wet conditions they can cause structural damage to buildings.
	Effects:	Common allergens and are also associated with hypersensitivity pneumonitis.
Bipolaris Drechslera	Habitat:	They are found in soil and as plant pathogens. Can grow indoors on a variety of substrates.
	Effects:	They may be allergenic and are very commonly involved in allergic fungal sinusitis. They are opportunistic pathogens but occasionally infect healthy individuals, causing keratitis, sinusitis and osteomyelitis.
Cladosporium	Habitat:	One of the most common genera worldwide. Found in soil and plant debris and on the leaf surfaces of living plants. The outdoor numbers are lower in the winter and often relatively high in the summer, especially in high humidity. The outdoor numbers often spike in the late afternoon and evening. Indoors, it can be found growing on textiles, wood, sheetrock, moist window sills and in HVAC supply ducts.
	Effects:	A common allergen, producing more than 10 allergenic antigens and a common cause of hypersensitivity pneumonitis.



1818 New York Ave. Suite 217 Washington, DC, 20002 (443) 691-0455

BB203 PGCPS Indoor Air Quality Beacon Heights Elementary School

#20044834

Organism Descriptions

Epicoccum

It is found in soil and plant litter and is a plant pathogen. It can grow indoors on a variety of substrates, including paper and textiles and is


commonly found on wet drywall.

It is a common allergen. No cases of infection have been reported in humans.

Myxomycetes

Found on decaying plant material and as a plant pathogen. Habitat:

Effects: Some allergenic properties reported, but generally pose no health concerns to humans.





Job Number: BB203

Company: Globa

Address: 1818 New York Avenue.

Beacon Heights Elementary School

Job Name: pGCPS Indoor Air Quality-

SHIP: FEDEX - PAK 50 DATE: 12-01-2020

MOLD

8160 4411 5679

Mobile: 443-691-0455 Email: channab@globalincusa.net

T	Collector: Kenna Leonzo			Beacon Heights Elementary School			ile: 443-69	1-0455	Email: channab@globalincusa.net			
_	Date Collected: 11/30/2020				; ;			Note:				
-	-	Analysis Type			Analysis Description			urnaround	Accepted Media Types			
-	Spore Trap S Identification 8				on & Enumeration of Fungal Spores	24	Hour	Air Cassettes, Impact Slides Air Cassettes, Impact Slides				
					Analysis with Dander, Fiber, and Pollen counts	24	Hour					
-	Direct	: ID	D	ID & Semi-Q	Quantative Enumeration of spores and mycelium		24	Hour	Bio-Tape, Tape, Swab, Bulk, Agar Plate			
-					ysis with Fully Quantitative spore count	24	Hour	Bio-Tape,	Bio-Tape, Tape, Swab, Bulk, Agar Plate			
	Cultur	re	C1	Identification	on & Enumeration of Mold only	7 0	ay	Air Plate,	Air Plate, Agar Plate, Swab, Bulk			
					on & Enumeration of Bacteria only	4 0	ay	Air Plate, Agar Plate, Swab, Bulk				
H			C3	Identification	on & Enumeration of Mold and Bacteria		7.0	ay	Air Plate, Agar Plate, Swab, Bulk			
			C5	Coliform Sc	creen for Sewage Bacteria		2 [ay	Agar Plate, Swab, Bulk			
				Total Partic	culate Analysis, ID & Count (Does Not Include Mole	d)	24	Hour	Air Cassettes, Impact Slides, Bio-Tape			
-	# Number		ber		Sample	Analy	sis	Volume		Notes		
o	1	BHES-1130-01 Ambie			nt	Ş	3	75 L				
	2			Room 1	19							
	3 BHES-1130-03		Room 9	9								
11	4	BHES-	1130-04	Multipu	ırpose		/	V				
十	5											
	6											
-	7						10,010					
-	8											
	9											
	10											
Ì	11	11										
	12											
	14											
	15	5										
-		 				i			1			

Released by: Kenna Leonzo

Date: 11/30/20

Received By:

12/

Date:

16